Author Index

Acheson, C. M., see Coppella, S. J. 189
Acree, Jr., W. E.
Thermochemical investigations of gas–liquid chromatography. IV. Partition coefficients of solutes on binary mixtures containing two complexing solvents 41
Anazawa, N., see Jinno, K. 173
Ballester, P., see García-Raso, A. 323
Bergueiro, R., see García-Raso, A. 323
Biggs, W. R., see Jinno, K. 173
Birsner, U.
——, Gilles, U., Nielsen, P. and McMaster, G. K. Reversed-phase high-performance liquid chromatographic system for the rapid, automated purification of oligonucleotides 381
Boer, G. de, see Kubo, I. 354
Burger, B. V.
—— and Munro, Z.
Headspace gas analysis. Liquid desorption of headspace volatiles trapped on activated carbon open tubular traps 95
Casella, M. L., see Castells, R. C. 65
Castells, R. C.
—— and Casella, M. L.
Evaporation rates of solvent blends measured by gas chromatography 65
Catterick, T., see White, P. C. 135
Chan, L. T. F.
——, Chhuy, L. H. and Crowley, R. J.
Gas chromatographic analysis of verapamil and norverapamil in *post mortem* specimens using a 530-μm fused-silica column with nitrogen–phosphorus detection 361
Cheetham, P. S. J., see Cookson, D. 265
Chhuy, L. H., see Chan, L. T. F. 361
Collins, C. H., see Valente, A. L. P. 349
Cookson, D.
——, Cheetham, P. S. J. and Rathbone, E. B.
Preparative high-performance liquid chromatographic purification and structural determination of 1-O-α-D-glucopyranosyl-D-fructose (trehalulose) 265
Coppella, S. J.
——, Acheson, C. M. and Dhurjati, P.
Isolation of high-molecular-weight nucleic acids for copy number analysis using high-performance liquid chromatograph 189
Cowman, M. K., see Hittner, D. M. 149
Crowley, R. J., see Chan, L. T. F. 361
D'Agostino, P. A.
——, Provost, L. R. and Visentini, J.
Analysis of O-ethyl S-[2-diisopropylamino)-ethyl] methylphosphonothiolate (VX) by capillary column gas chromatography–mass spectrometry 221
De Boer, G., see Kubo, I. 354
Dhurjati, P., see Coppella, S. J. 189
Dimitrov, Chr., see Gavrilova, T. B. 87
Femia, R. A.
—— and Weinberger, R.
Determination of reducing and non-reducing carbohydrates in food products by liquid chromatography with post-column catalytic hydrolysis and derivatization. Comparison with refractive index detection 127
Fernández-Sánchez, E.
——, García-Domínquez, J. A., Menéndez, V., Santiuste, J. M. and Pertierra-Rjmada, E.
Reduction of the number of characterizing probes in gas chromatography by factor analysis 318
Fetzer, J. C., see Jinno, K. 173
Flann, B.
—— and Lodge, B.
Analysis of estrogen sulphate mixtures in pharmaceutical formulations by reversed-phase chromatography 273
Fujimoto, Y.
——, Ishi, K., Nishi, H., Tsumagari, N., Kakimoto, T. and Shimizu, R.
New derivatization reagents for the resolution of carboxylic acid enantiomers by high-performance liquid chromatography 344
Fukuda, H., see Ohta, M. 392
García-Domínguez, J. A., see Fernández-Sánchez, E. 318
García-Raso, A.
——, Ballester, P., Bergueiro, R., Martínez, I., Sanz, J. and Jimeno, M. L.
Estimation of the polarity of stationary phases by proton nuclear magnetic resonance spectroscopy. Application to phenyl and methyl silicones (OV and SE series) 323
Gavrilova, T. B.
——, Parshina, I. V., Dimitrov, Chr., Ivanov, St., Petsev, N. and Topalova, I.
Carbon black adsorbents modified with *n*-hydrocarbons and treated with an high-frequency low-temperature plasma 87
Gelfi, C., see Rabilloud, T. 105
Gilles, U., see Birsner, U. 381

Goca, B., see Witkiewicz, Z. 73
Griffith, P. E., see Robinson, J. L. 201
Hadžija, O.
——, Iskrić, S. and Tonković, M.
Mobility of Fe(III)–phenolic acid complexes in thin-layer chromatography 358
Hagemeier, C. J., see Martin, C. F. 376
Hall, L. K., see Martin, C. F. 376
Hashimoto, Y., see Takeuchi, T. 328
Hittner, D. M.
—— and Cowman, M. K.
High-performance gel permeation chromatography of glycosaminoglycans. Column calibration by gel electrophoresis 149
Hogstrand, C., see Olsson, P.-E. 293
Horie, M.
——, Saito, K., Hoshino, Y., Nose, N., Mochizuki, E. and Nakazawa, H.
Simultaneous determination of nalidixic acid, oxolinic acid and piromidic acid in fish by high-performance liquid chromatography with fluorescence and UV detection 301
Hoshino, Y., see Horie, M. 301
Ishi, K., see Fujimoto, Y. 344
Ishii, D., see Takeuchi, T. 328
Iskrić, S., see Hadžija, O. 358
Itabashi, Y.
—— and Takagi, T.
High-performance liquid chromatographic separation of diacylglycerol enantiomers on a chiral stationary phase 257
Ivanov, St., see Gavrilova, T. B. 87
Iwaki, K., see Nimura, N. 387
Jimeno, M.L., see García-Raso, A. 323
Jinno, K.
——, Anazawa, N., Okamoto, M., Tanaka, N., Fetzer, J. C. and Biggs, W. R.
Elution behaviour of peropyrene-type polycyclic aromatic hydrocarbons in liquid chromatography 173
John, J., see Robinson, J. L. 201
Kakimoto, T., see Fujimoto, Y. 344
Kasai, M., see Sawada, S. 283
Kim, M., see Kubo, I. 354
Kimura, T., see Ohta, M. 392
Kinoshita, T., see Nimura, N. 387
Kirkes, K. A., see Robinson, J. L. 201
Klesper, E.
—— and Schmitz, F. P.
Single and multiple gradients in supercritical fluid chromatography (Review) 1
Kubo, I.
——, Kim, M. and De Boer, G.
Efficient isolation of insect growth inhibitory macrolide alkaloids using recycle high-performance gel permeation chromatography 354
Kulanthaivel, P.
—— and Pelletier, S. W.
High-performance liquid chromatographic analysis of C_{19}-diterpenoid alkaloids 366
Lambert, N.
——, Plumb, G. W. and Wright, D. J.
Application of high-performance liquid chromatography to the assessment of subunit heterogeneity in plant 11S storage globulins 159
Lau, B. P.-Y.
——, Weber, D. and Page, B. D.
Gas chromatographic–mass spectrometric determination of ethyl carbamate in alcoholic beverages 233
Lodge, B., see Flann, B. 273
McMaster, G. K., see Birsner, U. 381
Martin, C. F.
——, Takahashi, L. T., Worsley, J. L., Hagemeier, C. J. and Hall, L. K.
Analysis of sodium cefazolin in ophthalmic ointment by high-performance liquid chromatography 376
Martínez, I., see García-Raso, A. 323
Masuda, H., see Mihara, S. 309
Menéndez, V., see Fernández-Sánchez, E. 318
Mihara, S.
—— and Masuda, H.
Correlation between molecular structures and retention indices of pyrazines 309
Mochizuki, E., see Horie, M. 301
Munro, Z., see Burger, B. V. 95
Nakazawa, H., see Horie, M. 301
Nicoll-Griffith, D. A.
Stereoelectronic model to explain the resolution of enantiomeric ibuprofen amides on the Pirkle chiral stationary phase 179
Nielsen, P., see Birsner, U. 381
Nimura, N.
——, Iwaki, K. and Kinoshita, T
Separation of norepinephrine enantiomers, DOPA enantiomers and dopamine derivatized with *o*-phthalaldehyde–N-acetyl-L-cysteine by high-performance liquid chromatography 387
Nishi, H., see Fujimoto, Y. 344
Nonhebel, H. M.
Direct separation of (*S*)- and (*R*)-abscisic acid on a commercially available chiral high-performance liquid chromatographic column 374
Nose, N., see Horie, M. 301
Ohta, M.
——, Fukuda, H., Kimura, T. and Tanaka, A.
Quantitative analysis of oxytocin in pharmaceutical preparations by high-performance liquid chromatography 392

Okamoto, M., see Jinno, K. 173
Olsson, P.-E.
—— and Hogstrand, C.
Improved separation of perch liver metallothionein by fast protein liquid chromatography 293
Page, B. D., see Lau, B. P.-Y. 233
Papp, E.
System peaks in ion chromatography using cetrimide-coated columns and indirect photometric detection 211
Parshina, I. V., see Gavrilova, T. B. 87
Pelletier, S. W., see Kulanthaivel, P. 366
Pernelle, J.-J., see Rabilloud, T. 105
Pertierra-Rimada, E., see Fernández-Sánchez, E. 318
Petsev, N., see Gavrilova, T. B. 87
Plumb, G. W., see Lambert, N. 159
Provost, L. R., see D'Agostino, P. A. 221
Rabilloud, T.
——, Pernelle, J.-J., Wahrmann, J. P., Gelfi, C. and Righetti, P. G.
Removal of oligomers and *n*-mers from the Immobiline chemicals for isoelectric focusing 105
Railton, I. D.
Resolution of the enantiomers of abscisic acid methyl ester by high-performance liquid chromatography using a stationary phase of cellulose tris(3,5-dimethylphenylcarbamate)-coated silica gel 371
Rashid, Z. A., see Szymanowski, J. 55
Rathbone, E. B., see Cookson, D. 265
Ravichandran, K.
—— and Rogers, L. B.
Chiral separations of amino acids using di- and tri-peptides for zwitter-ion pair chromatography 49
Richards, M. P.
—— and Steele, N. C.
Isolation and quantitation of metallothionein isoforms using reversed-phase high-performance liquid chromatography 243
Righetti, P. G., see Rabilloud, T. 105
Robinson, J. L.
——, John, J., Safa, A. I., Kirkes, K. A. and Griffith, P. E.
Evaluation of disposable cartridges for trace enrichment from aqueous solutions 201
Rogers, L. B., see Ravichandran, K. 49
Safa, A. I., see Robinson, J. L. 201
Saito, K., see Horie, M. 301
Santiuste, J. M., see Fernández-Sánchez, E. 318
Sanz, J., see García-Raso, A. 323
Sawada, S.
——, Kasai, M., Yamada, T. and Takahashi, T.
Determination of photosynthetic intermediates in leaves by high-performance liquid chromatography 283
Schmitz, F. P., see Klesper, E. 1
Shimizu, R., see Fujimoto, Y. 344
Steele, N. C., see Richards, M. P. 243
Szymanowski, J.
——, Voelkel, A. and Rashid, Z. A.
Polarity and hydrophile–lipophile balance of model hydroxyoximes and their intermediates as determined by reversed-phase gas chromatography 55
Takagi, T., see Itabashi, Y. 257
Takahashi, L. T., see Martin, C. F. 376
Takahashi, T., see Sawada, S. 283
Takeuchi, T.
——, Hashimoto, Y. and Ishii, D.
Temperature effect in packed microcolumn chromatography using a carbon dioxide mobile phase 328
Tanaka, A., see Ohta, M. 392
Tanaka, N., see Jinno, K. 173
Theis, D. L.
Ion-pairing liquid chromatographic method for the determination of pirlimycin hydrochloride 335
Tonković, M., see Hadžija, O. 358
Topalova, I., see Gavrilova, T. B. 87
Tsumagari, N., see Fujimoto, Y. 344
Valente, A. L. P.
—— and Collins, C. H.
Gas chromatographic quantification of bromine-containing compounds using a flame ionization detector 349
Visentini, J., see D'Agostino, P. A. 221
Voelkel, A., see Szymanowski, J. 55
Wahrmann, J. P., see Rabilloud, T. 105
Weber, D., see Lau, B. P.-Y. 233
Weinberger, R., see Femia, R. A. 127
Wheals, B. B.
Ion chromatography of inorganic anions on a dynamically modified polystyrene–divinylbenzene packing material and its application to anion screening by sequential ultraviolet absorbance and electrochemical detection 115
White, P. C.
—— and Catterick, T.
Evaluation of absorbance ratioing for solute identification in high-performance liquid chromatography using a diode array detector 135

Witkiewicz, Z.
—— and Goca, B.
Disc-like liquid crystal stationary phases derived from hexahydroxybenzene 73

Worsley, J. L., see Martin, C. F. 376
Wright, D. J., see Lambert, N. 159
Yamada, T., see Sawada, S. 283

Erratum

J. Chromatogr., 395 (1987) 523–529
Page 525, last line, "30–15%" should read "3–15%".